

**COVID-19 Content**

# What Is the Preparedness and Capacity of Palliative Care Services in Middle-Eastern and North African Countries to Respond to COVID-19? A Rapid Survey



Sabah Boufkhed, Richard Harding, Tezer Kutluk, Abdullatif Husseini, Nasim Pourghazian, and Omar Shamieh

*Cicely Saunders Institute (S.B., R.H.), Florence Nightingale Faculty of Nursing Midwifery and Palliative Care, King's College London, London, United Kingdom; Department of Pediatric Oncology Hacettepe University Faculty of Medicine and Cancer Institute (T.K.), Ankara, Turkey; Institute of Community and Public Health (A.H.), Birzeit University, Birzeit, Palestine; Regional Office for the Eastern Mediterranean (N.P.), World Health Organization, Cairo, Egypt; Center for Palliative & Cancer Care in Conflict (O.S.), Department of Palliative Care, King Hussein Cancer Center, Amman; and College of Medicine (O.S.), The University of Jordan, Amman, Jordan*

**Abstract**

**Context.** Evidence from prior public health emergencies demonstrates palliative care's importance to manage symptoms, make advance care plans, and improve end-of-life outcomes.

**Objective.** To evaluate the preparedness and capacity of palliative care services in the Middle-East and North Africa region to respond to the COVID-19 pandemic.

**Methods.** A cross-sectional online survey was undertaken, with items addressing the WHO International Health Regulations. Nonprobabilistic sampling was used, and descriptive analyses were conducted.

**Results.** Responses from 43 services in 12 countries were analyzed. Half of respondents were doctors (53%), and services were predominantly hospital based (84%). All but one services had modified at least one procedure to respond to COVID-19. Do Not Resuscitate policies were modified by a third (30%) and unavailable for a fifth (23%). While handwashing facilities at points of entry were available (98%), a third had concerns over accessing disinfectant products (37%), soap (35%), or running water (33%). The majority had capacity to use technology to provide remote care (86%) and contact lists of patients and staff (93%), though only two-fifths had relatives' details (37%). Respondents reported high staff anxiety about becoming infected themselves (median score 8 on 1–10 scale), but only half of services had a stress management procedure (53%). Three-fifths had plans to support triaging COVID-19 patients (60%) and protocols to share (58%).

**Conclusion.** Participating services have prepared to respond to COVID-19, but their capacity to respond may be limited by lack of staff support and resources. We propose recommendations to improve service preparedness and relieve unnecessary suffering. *J Pain Symptom Manage* 2021;61:e13–e50. © 2020 Published by Elsevier Inc. on behalf of American Academy of Hospice and Palliative Medicine.

**Key Words**

*Palliative care, preparedness, COVID-19, pandemic, epidemic, Middle-East and North Africa*

Authors' contributions: SB and RH developed the original protocol; SB, RH, OS, TK adapted the survey questionnaire and protocol; SB, RH, OS, TK, AH and NP commented on the methods and conducted data collection; SB conducted the analysis and drafted the manuscript; SB, RH, OS, TK, AH and NP interpreted the findings. All authors revised and approved the final draft.

*Address correspondence to:* Sabah Boufkhed, Florence

Nightingale Faculty of Nursing Midwifery and Palliative Care, Cicely Saunders Institute, Department of Palliative Care, Policy & Rehabilitation, King's College London, Bessemer Road, London SE5 9RS, United Kingdom. E-mail: [sabah.boufkhed@kcl.ac.uk](mailto:sabah.boufkhed@kcl.ac.uk)

*Accepted for publication:* October 23, 2020.

## Key message

This study addresses an important gap in the preparedness of palliative care services in the Middle-East and North Africa region to respond to COVID-19 pandemic and other outbreaks. Our survey led to seven recommendations aiming at improving their preparedness and supporting the region's health systems in relieving unnecessary suffering.

## Introduction

Elderly people and those with underlying health conditions, such as cancer, are most at risk of developing severe COVID-19 or dying.<sup>1–3</sup> Comorbidities such as diabetes and cardiovascular disease are highly prevalent in the Middle-East, raising concerns for the progress of the pandemic in the region.<sup>4</sup> In weaker health systems, there is limited capacity to care for COVID-19 patients who require intensive care units because of moderate to severe forms of the disease or complex symptoms such as breathlessness.<sup>2,5</sup>

Palliative care is explicitly recognized under the human right to health, relieving the suffering of patients and families and improving their quality of life and outcomes while saving health-care costs.<sup>6–8</sup> The World Health Assembly has called on countries to provide palliative care in the clinical management of COVID-19 patients.<sup>9,10</sup> This would address the emerging evidence for palliative care needs among COVID-19 patients, including physical symptoms (e.g. fever, breathlessness, fatigue, cough),<sup>2,11</sup> spiritual distress related to survival, and psychological distress related to prognosis uncertainty.<sup>12,13</sup>

Under the 2015 International Health Regulations (IHR), countries are required to prepare response plans for public health emergencies of international concern.<sup>14,15</sup> However, preparedness plans routinely fail to include palliative care.<sup>16–18</sup> The role of palliative care in pandemic responses has been demonstrated, including sharing symptom management protocols and training nonpalliative care health-care workers, supporting patients' triage and providing psychosocial and bereavement care.<sup>10</sup> The "COVID-19 tsunami of suffering" is likely to increase the need for palliative care, especially in low- and middle-income countries.<sup>8</sup>

In 2012, national health systems in the Middle East region used their influenza surveillance systems to detect the MERS-CoV, and most countries have tested their preparedness plans.<sup>14,19–25</sup> In Middle Eastern countries with fragile or limited health-care systems, the COVID-19 pandemic may cause enormous challenges to fragile health systems.<sup>26,27</sup>

Palliative care in the Middle East and North African region is a relatively new development, with no

country having fully integrated it within the health system.<sup>28,29</sup> With a Muslim majority in the region, the pandemic may have an additional religious or spiritual impact for populations.<sup>30–33</sup> A systematic review of end-of-life care in Muslim-majority countries highlighted the central role of families in the decision-making process, as well as the need for spaces to perform rituals and to address preferences for pain management.<sup>34</sup> The preparedness and capacity of palliative care services to respond to COVID-19 in the MENA region is still unknown. To inform policy and appropriate and timely responses, we aimed to evaluate the preparedness and capacity to respond to COVID-19 within palliative care services of the Middle-East and North Africa region.

## Method

### Study Design and Settings

We developed, piloted, and conducted an exploratory cross-sectional online survey, using WHO's 2005 International Health Regulations<sup>25</sup> and online survey methodological guidelines.<sup>35,36</sup>

### Population and Sampling

Nonprobabilistic sampling combining convenience and snowball sampling was used to recruit representatives of palliative care services in the Middle-East and North Africa region: Afghanistan, Algeria, Bahrain, Egypt, Iran, Iraq, Jordan, Kuwait, Lebanon, Libya, Morocco, Oman, Palestine, Pakistan, Qatar, Saudi Arabia, Syria, Tunisia, Turkey, United Arab Emirates, Yemen. We aimed to recruit at least one service per country and to have only one respondent per service. The services included hospices, hospital services, and home- and community-based care.

There is no comprehensive formal list, registry, or network of palliative care services for the Middle-East region. There are limited palliative care services and publicly available information on the existing official registry of palliative care services for the countries that could give a framework to draw a representative sample of services. Therefore, we identified services using publicly available information and the research team's networks as follows. We yielded 60 contacts from the Atlases of Palliative Care,<sup>37–39</sup> International Association for Hospices and Palliative Care (IAHPC) directory,<sup>40</sup> and a rapid Google and PubMed search. We yielded a further 160 contacts from our professional networks, partners of Research For Health in Conflict in the Middle-East and North Africa ([r4hcmena.org](http://r4hcmena.org)), and disseminated our survey through the World Health Organization's Eastern Mediterranean Region (WHO-EMRO) network for palliative care. Countries from WHO-EMRO located in sub-Saharan

Africa were approached in a separate survey we conducted with the African Palliative Care Association.<sup>41</sup>

### Data Collection

The original survey questionnaire was initially designed by researchers from Italy and the UK for an early assessment of the Italian palliative care response early during the COVID-19 pandemic who shared with us the questionnaire in English.<sup>42</sup> We further developed it to include existing international recommendations, especially the IHR, and evidence for generic preparedness to respond to infectious disease outbreaks or pandemics,<sup>19,23–25,43–47</sup> and recommendations for palliative care response and roles in epidemics and pandemics.<sup>16,17</sup> R4HC-MENA research team members adapted the survey to the region using professional experience and key preparedness literature for the region.<sup>19,20,26,48</sup> The full questionnaire was available in English and Arabic (see Appendix I and II). It was developed in English then translated into Arabic by a bilingual professional translator, and the translations were reviewed by two independent experts. The translation was then tested during the pilot, where two researchers not involved in the translation completed both versions and gave feedback. No corrections were suggested.

Data were collected online using the *SmartSurvey*<sup>TM</sup> platform, permitting one answer per computer on the online platform, and enabled the anonymity function so IP addresses were not collected. The survey link was emailed to 190 contacts across the region, inviting them to participate (institutions providing care) and disseminate (via professional networks).

Recruitment to the online survey was opened on 05/26/20 and closed on 06/22/2020, with responses permitted until 06/26/2020. Reminders to complete the survey were sent twice during the period.

### Data Analysis

Descriptive analyses were conducted following de-duplication. Quantitative data were analyzed using Stata (version 16): categorical variables described as frequencies and percentage; continuous variables as median and interquartile range (IQR). Open-ended questions were coded thematically, and the dominant themes reported.<sup>49</sup>

### Ethics

Ethical approval was obtained from King's College London Research Ethics Office (reference LRS-19/20–19,091). Data were collected and stored following the UK 2016 General Data Protection Regulation. Informed consent was obtained online from participants within the survey.

## Findings

### Respondents and COVID-19 Situation in Their Services

There were 113 engagements with the online survey (65 for the English version and 48 for the Arabic one), 69 individuals completed the survey (37 in English and 32 in Arabic) of which 67 gave consent (completion rate: 59%). The final sample was 43 (33 English and 10 Arabic) after de-duplication. Of 21 countries in the region, 12 had a least one participant (57%). We received no participation from nine countries (Afghanistan, Algeria, Iraq, Qatar, Tunisia, United Arab Emirates, Yemen) (33%). For two countries (Libya and Syria), we could not identify a contact (9%).

Table 1 describes the participants' characteristics. Half were medical doctors (53%) and one-fifth were nurses (19%). Two in five services were public or governmental (40%) and the majority were hospital based (84%). Responding services had a median of 500 patients per year (IQR: 200–2500; 4 missing data), and 33 services reported having beds (median: 13; IQR: 8–25).

Table 1  
Respondents' Characteristics (N = 43)

	n	%
Country		
Turkey	14	33
Jordan	12	28
Other (Bahrain, Egypt, Iran, Kuwait, Lebanon, Morocco, Oman, Pakistan, Palestine, Saudi Arabia: 1 to 3 respondents per country)	17	40
Respondent's current role(s)		
Doctor or medical officer	23	53
Nurse	8	19
Manager or responsible for the service and doctor, nurse or psychosocial professional <sup>a</sup>	5	12
Manager or responsible for the service	3	7
Psychosocial professional	3	7
Other (Operations Manager)	1	2
Type of organisation		
Public	17	40
Nonprofit charity	5	12
Mixed <sup>b</sup>	4	9
Private	3	7
Missing	14	33
Type of service <sup>c</sup>		
Within hospital	36	84
Within community	13	30
Outpatient	10	23
Inpatient	7	16
Hospice/service having beds	33	77
Services which reported a case (possible, suspect or confirmed)	27	63

<sup>a</sup>Doctor + Manager or responsible of the service (n = 3); Nurse + Manager or responsible of the service (n = 1); Nurse + Psychosocial professional + Manager or responsible of the service (n = 1).

<sup>b</sup>Mixed: public + nonprofit (n = 2); public + private (n = 1).

<sup>c</sup>Multiple choices allowed.

At the time of the survey, two-thirds of respondents had experienced a COVID-19 case within their service, with a median of five cases (IQR: 3–10.5) (see [Appendix III](#) and [IV](#)). Most cases included patients (81%) and were identified in another service within the hospital (61%).

### *Policies and Procedures*

[Table 2](#) describes the policies and procedures available or modified in response to the COVID-19 pandemic. While the vast majority reported having a written procedure for “what to do” in case of COVID-19 in the service among patients, staff, volunteers (77–88%), a third did not have or were unsure (21% and 9% respectively) to have a COVID-19 case definition (confirmed, probable, and suspect).

All but one service had modified at least one of their procedures in response to the COVID-19 pandemic, especially for operators’ protection (91%) and visitors and relatives (88%). However, fewer had modified their volunteer support (56%) and Do Not Resuscitate (30%) policies, with one-quarter not having these policies in place (i.e. N/A response; 23%).

### *Infection Control Measures and Resources*

The majority of participants reported having in place several measures to control infection and knew how to access resources in case of outbreak or

lockdown (see [Tables 3](#) and [4](#)). All but one had hand-washing facilities at entry points and half of those were there before COVID-19. Four in five provided additional personal protective equipment for palliative care staff (84%) and cleaners (79%) and identified an isolation room (81%). A majority had up-to-date inventories of medicines and medical supplies (86%) and personal protective equipment (81%) available. Respondents knew how to dispose of highly infectious waste in the service (84%), had materials and facilities to dispose of it (91%), and had staff trained in handling highly infectious conditions (90%), of which half were trained before COVID-19. However, one-third had concerns regarding access to essential resources for ensuring safe care: disinfectant products (37%), soap (35%), hand sanitizers (35%), running water (33%), contactless thermometers (33%), and electricity (28%). These concerns were higher for access in the surrounding community.

### *Information Systems, Communication, and Technology*

The respondents had communication channels identified for use during the pandemic (see [Appendix V](#)). Three in four had a designated focal point for collecting and sharing up-to-date information (72%). The majority would use mobile phones to receive information (77%) and phone calls to share

*Table 2*  
**Procedures (or Guidance) in Place and Policies Modified (N = 43)**

	Yes	No	Unsure or Do not Know	Missing or N/A
	n (%)	n (%)	n (%)	n (%)
Case definition for confirmed, probable, and suspected COVID-19 cases	30 (70)	9 (21)	4 (9)	0 (0)
Written procedure for “what to do” in the service in case of COVID-19 case among				
Patients	38 (88)	4 (9)	0 (0)	1 (2)
Health-care professional staff member	38 (88)	3 (7)	1 (2)	1 (2)
Volunteers and medical staff	35 (81)	5 (12)	2 (5)	1 (2)
Relatives and visitors	33 (77)	6 (14)	3 (7)	1 (2)
Staff and volunteers going in the community	28 (65)	7 (16)	7 (16)	1 (2)
Written procedure for “what to do” in the service in case of infectious diseases among				
Patients	28 (65)	6 (14)	5 (12)	4 (9)
Relatives and visitors	24 (56)	7 (16)	7 (16)	5 (12)
Health-care professional staff member	30 (70)	4 (9)	6 (14)	3 (7)
Volunteers and medical staff	26 (60)	5 (12)	7 (16)	5 (12)
Staff and volunteers going in the community	19 (44)	8 (19)	12 (28)	4 (9)
Policies or procedures modified as a measure to avoid contagion				
Operators’ protection (personal protective equipment)	39 (91)	1 (2)	0 (0)	3 (7)
Visitors/relatives (number of visitors, hours, etc.)	38 (88)	0 (0)	2 (5)	3 (7)
Dead body handling	35 (81)	4 (9)	3 (7)	1 (2)
Patients’ admission to the service	34 (79)	1 (2)	3 (7)	5 (12)
Volunteer support	24 (56)	6 (14)	6 (14)	7 (16)
Care of the relatives after the patient’s death	19 (44)	13 (30)	7 (16)	4 (9)
Do Not Resuscitate	13 (30)	16 (37)	4 (9)	10 (23)
Procedure to support health-care providers to manage stress	23 (53)	16 (37)	4 (9)	0 (0)
Recommendations if you or someone in your household becomes ill with COVID-19 symptoms	39 (91)	3 (7)	1 (2)	0 (0)

Table 3  
Measures Taken to Avoid Contagion (N = 43)

	Additional Ones/Trained Because of COVID-19		Already Before COVID-19		None	
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Hand washing facility for all at points of entry	22 (51)		20 (47)		1 (2)	
Personal protection equipment (PPE) for						
Palliative care staff	36 (84)		6 (14)		1 (2)	
Cleaning staff <sup>a</sup>	34 (79)		7 (16)		1 (2)	
All health-care providers have been trained in handling highly infectious conditions such as COVID-19	16 (37)		23 (53)		4 (9)	

<sup>a</sup>1 missing data.

information in case of emergency (65% with staff; 91% with patients and relatives).

Almost all services had up-to-date lists of patients and staff (93%), and collected patients' symptoms, outcomes (95%) and treatment (98%) (see Appendix VI). Two-thirds used electronic records to collect the latter health information (81%). However, two in five services did not collect visitors' and relatives' contact details or visit dates (37%).

The vast majority reported having the capacity to use technology to provide remote care (86%). Twenty-seven participants shared perceived advantages and disadvantages of using technology (see Appendix VII–XII). The key limitations were a lack of resources (e.g., lack of Internet coverage or devices for patients) ( $n = 9$ ); trust issues from users and cooperation from patients' family ( $n = 4$ ), appropriateness issues regarding age or condition ( $n = 3$ ) or a lack of

Table 4  
Resources Available and Access and Knowledge (N = 43)

	Yes	No	Do not Know/Not Sure	Missing
	n (%)	n (%)	n (%)	n (%)
Adequate material and facilities to dispose of highly infectious waste				
In the hospice	39 (91)	4 (9)	0	0
In the community	19 (44)	6 (14)	17 (40)	1 (2)
Up-to-date inventory of				
Protection material available for staff, patient, and visitors	35 (81)	2 (5)	2 (5)	0 (0)
Medicines and other medical supplies available	37 (86)	2 (5)	2 (5)	0 (0)
Capacity to use technology instead of face-to-face appointment to provide some care remotely	37 (86)	6 (14)	—	—
Phone call	34 (92)	3 (8)	—	—
Video call	20 (54)	17 (46)	—	—
Concerns about the service/hospice's access to				
Disinfectant products	16 (37)	26 (60)	—	1 (2)
Soap	15 (35)	25 (58)	—	3 (7)
Hand sanitizers (with 60% alcohol)	15 (35)	27 (63)	—	1 (2)
Running water	14 (33)	28 (65)	—	1 (2)
Thermometers (contactless, thermoflash-type)	14 (33)	28 (65)	—	1 (2)
Electricity	12 (28)	30 (70)	—	1 (2)
Having concerns about the surrounding's access to				
Accessing disinfectant products to continue providing care safely	20 (47)	20 (47)	—	3 (7)
Hand sanitizers (with 60% alcohol)	19 (44)	21 (49)	—	3 (7)
Thermometers (contactless, thermoflash-type)	15 (35)	24 (56)	—	4 (9)
Soap	13 (30)	22 (51)	—	8 (19)
Running water	11 (26)	27 (63)	—	5 (12)
Electricity	9 (21)	27 (63)	—	7 (16)
Knowledge of how the hospice/service would access to the following in case of emergency, lockdown or quarantine				
Food (N = 36 - for hospital-based or inpatient services only)	31 (86)	11 (26)	—	1 (4)
Medicines and other medical supply	38 (88)	5 (12)	—	0 (0)
Additional staff (e.g. if staff self-isolates or becomes ill)	37 (86)	4 (9)	—	2 (5)
Knowledge of how to dispose of to dispose of highly infectious waste				
In the hospice or service	36 (84)	4 (9)	—	3 (7)
In the community	27 (63)	9 (21)	—	7 (16)
Cleaning staff included in information sharing and training regarding managing COVID-19	34 (79)	3 (7)	—	6 (14)
Having education material about COVID-19 available	36 (84)	6 (14)	—	1 (2)
Posters displayed where staff, patients, and visitors can see them (N = 36)	33 (92)	3 (8)	—	—
Education material also available for the surrounding community (N = 36)	31 (86)	1 (2)	—	1 (2)

body language to effectively communicate ( $n = 3$ ) were the most reported. One participant was concerned about relatives hiding information from the patients. However, several advantages were also shared, such as the ability to deliver care and manage patients remotely ( $n = 8$ ); and control of potential transmission ( $n = 6$ ).

### *Palliative Care Staff and Expertise to Support Pandemic Response*

Respondents reported high anxiety among staff, especially risk to them of becoming infected (median score 8 on 1–10 scale; IQR: 7–9) and their ability to care for their relatives (median score 8 on 1–10 scale; IQR: 6–9) (see Table 5). However, only half of the services had a staff stress management procedure (53%).

Table 6 shows that responding services have the capacity to support the broader health system to respond to COVID-19. Three in five services had plans to support other health-care settings in triaging COVID-19 patients (60%) and protocols to share for symptom management and psychological support (58%). Among the 25 respondents who could share those protocols, the majority declared they could train nonspecialists in using them (72%). About half of the services had redeployment plans for palliative care staff and resources, although 20% did not know if they had such plans. A third of the services did not have or did not know about plans to redeploy volunteers.

## **Discussion**

Our survey provides the first comprehensive assessment of the preparedness of palliative care services in the Middle-East and North African region to respond to a pandemic. Responding services have prepared to respond to COVID-19, but their capacity to respond may be limited by the lack of access to infection control basics and community-based services, especially in case of a lockdown. Lack of support to staff in managing their stress and anxiety is a major

concern. This is crucial to equip them to deliver sustained care to existing non-COVID and new COVID-19 patients and their families and to fulfill their potential in supporting the wider health service during the epidemic.

In line with findings from our recent survey of African palliative care services,<sup>41</sup> the respondents were aware of the communication channels to be used in case of emergency, had up-to-date lists of staff and patients that would facilitate contact tracing in case of an outbreak, and had modified policies regarding operators' protection and visits. However, among the MENA region's respondents, there was a higher proportion of services using electronic records, which would facilitate rapid contact tracing and patients' monitoring in case of an outbreak. The region has developed electronic medical records,<sup>50,51</sup> (particularly in hospitals) and most respondents were hospital based. However, it is important to note that our rapid survey may be biased toward most advanced services that are part of national or international networks. Further research would be needed to assess the penetration of electronic health information systems in more remote and nonhospital-based palliative care services.

In line with recommendations on the role of palliative care services in responding to epidemics,<sup>16</sup> most services had protocols for symptom management and bereavement to share and were ready to support COVID-19 patients triage. This may be because two-thirds already had a COVID-19 case in the hospital they were based in or had learned from previous experience with MERS-CoV in the region. The lack of plans to redeploy staff, or resources, or of stress management procedures for staff identified threaten staff well-being and sustainability of patient care. The establishment of a regional association collecting and sharing protocols and resources in the languages of the region may help address this gap.

Regarding using technology to avoid face-to-face interactions, while the majority of services reported they had the capacity, it will be important to further investigate some of the barriers identified in the survey.

Table 5  
Perceived Effects of COVID-19 on Staff and Risks for the Service ( $N = 43$ )

	Median (IQR)
Perceived effects on staff	
Anxious about getting infected themselves <sup>a</sup>	8 (7–9)
Anxious about the need to care for their own relatives <sup>a</sup>	8 (6–9)
Anxious about the need to care for their children who may not be at school <sup>a</sup>	8 (6–9)
Worried regarding potential issues for their interaction with the community if the service is known to manage a potential COVID-19 case <sup>a</sup>	7 (5–8)
Perception of the risks in the coming week <sup>b</sup>	
Hospice/palliative care staff are at risk of being infected by COVID-19	5.5 (4–7)
Hospice/palliative care service is at risk of closing because of an infection in the hospice or service	5 (2–7)

<sup>a</sup>2 missing data.

<sup>b</sup>1 missing data.

Table 6  
Palliative Care Expertise to Support the Broader Health System (N = 43)

	Yes	No	Do not Know or N/A	Missing
	n (%)	n (%)	n (%)	n (%)
Plans to redeploy at least one of the following outside of the inpatient settings, in case of outbreak COVID-19 or another highly infectious disease				
Health-care providers	21 (49)	10 (23)	8 (19)	4 (9)
Resources (material and supplies)	19 (44)	10 (23)	8 (19)	6 (14)
Volunteers	12 (28)	10 (23)	6 (14)	7 (16)
Plans to support other health-care services in the triage of patients in case of COVID-19 outbreak	26 (60)	15 (35)	- (-)	2 (5)
Palliative care protocols for symptom management and psychological support that could be shared with nonspecialist staff and/or COVID-19 response teams in other health-care facilities	25 (58)	17 (40)	- (-)	1 (2)
If yes, capacity to train nonspecialist in using these protocols (N = 25)	18 (72)	6 (24)	- (-)	1 (4)

The barriers reported by participants, such as lack of resources or trust issues, reflect those reported in the wider e-health implementation literature.<sup>52</sup> Further research is needed to assess access and reliability of the connectivity required for e-health in MENA countries, focusing on the specificities of the regional and cultural context and related to Muslim-majority countries.<sup>34</sup> Technology has advantages to reducing potential infections and solutions to reaching hard-to-reach groups need to be further explored,<sup>53</sup> especially to reach out to forcibly displaced populations in a region widely affected by conflicts.

### Strengths and Limitations

This rapid survey provides the first comprehensive assessment of the preparedness of palliative care services in the MENA region. We used a standardized questionnaire using international guidelines and standards adapted to the region, enabling international comparison. We have insights for 12 countries from the region with a broad geographical range. The lack of a comprehensive list of PC services in the region may have introduced a sampling bias, which may limit the generalizability of our findings, but we developed a plan to identify and include as many eligible participants as possible. It is noteworthy that the limited number of respondents per country may also demonstrate the difficulty in contacting and mobilizing palliative care services in the region in response to public health emergencies of international concern. The questions regarding the availability of medicines that we used were generic as we used the IHR.<sup>25</sup> While detailed investigation into the challenges of drug availability was beyond the scope of this study, further investigation of the potential issues in accessing opioid, especially in MENA countries, is warranted. Finally, while having more than one respondent per service would have allowed capturing potentially different views from various staff, our

criteria of sampling the individual responsible for the service gave a rapid assessment in an urgent context. Although we de-duplicated responses based on key characteristics, more than one participant may have responded from a single service.

### Recommendations

This study provides urgently needed primary evidence to inform policy and practice in the region. Of utmost importance, we call for appropriate resources to support staff and palliative care services. We propose the following recommendations for policy and practice.

1. Governments and services should ensure that basic water and sanitation are available to ensure a safe provision of palliative care with implementation of infection control measures.
2. Governments and services should also allocate funding to equip their palliative care facilities, staff, and potentially the patients' access to devices, such as mobile smartphone.<sup>53</sup>
3. All palliative care services need to acquire stress management protocols and offer services to support the staff. Palliative care staff well-being and views should be assessed, especially before deciding to redeploy them.
4. Palliative care services need to be involved in supporting and training nonspecialist health-care workers in complex questions related to the care of the dying. This is particularly relevant in light of the scarcity of resources faced by the health systems and health-care professionals, raising ethical issues and difficult discussions related to the triage and resuscitation.<sup>54</sup>
5. We advise palliative care services in the region to develop DNR policies when absent and adapt them early before the crisis emerge.<sup>55</sup> Such decisions need to be evaluated using a social justice lens and inform ethical discussions on resuscitation debates on COVID-19.<sup>56,57</sup>

Palliative care services would need to ethically allocate their resources while maintaining dying patients' dignity, and ensure appropriate communication with caregivers.

6. We call on governments to integrate palliative care into the preparedness plans as recommended in the WHO publications, to prevent unnecessary suffering and foster a rapid and flexible response in case of public health emergencies. We also call on the WHO to revise the IHR to support and evaluate countries' preparedness progress and reflect the necessity of palliative care into the emergency.<sup>16</sup>
7. Finally, we propose the development of the National Palliative Care Reference Center in the MENA region based on the model of National Reference Laboratories in the WHO-IHR.<sup>15</sup> Such centers could be rapidly mobilized and foster the achievement of Universal Health Coverage. The National Palliative Care Reference Center could collect, compile, and share the most up-to-date information and protocols with other palliative care services within their countries and the region; but also train non-specialists and less advanced or resourced palliative care providers in case of emergency, and beyond. They could be coordinated by a regional association or an existing network such as the WHO-EMRO palliative care expert network.

### ***Disclosures and Acknowledgments***

The authors have no conflicts of interest to disclose. The authors would like to thank Shayma'a Turki, Ayman Issa and Dr. Nour Horanieh for their help in English-Arabic translations; Ghadeer Al-Arja and Waleed Alrjoob for their help in piloting the survey; and the World Health Organization's Eastern Mediterranean Region (WHO-EMRO) network for palliative care for their support and help in data collection.

Sabah Boufkhed, Richard Harding (RH), Tezer Kutluk, and Omar Shamieh are funded through the UK Research and Innovation GCRF Research for Health in Conflict (R4HC-MENA); developing capability, partnerships and research in the Middle and Near East (MENA) ES/P010962/1. The overall vision for the R4HC-MENA partnership is to build sustainable research and policy capacity in the region to address major health challenges arising from conflict. R4HC-MENA activities aim to facilitate more effective translation of research into policy and deliver impact on both the research community and for patients and vulnerable populations. Principal and Co-Investigators: Adam Coutts, Brendan Burchell, Cengiz Kılıç,

Deborah Mukherji, Fouad M Fouad, Ghassan Abu Sitah, Hanna Kienzler, Kai Ruggeri, Kalipso Chalkidou, Matthew Moran, Omar Shamieh, Preeti Patel, Richard Harding, Richard Sullivan, Rita Giacaman, Şevkat Bahar Özvarış, Simon Deakin, Tezer Kutluk, Wyn Bowen. RH is funded by the National Institute of Health Research (NIHR) Global Health Research Unit on Health System Strengthening in Sub-Saharan Africa, King's College London (GHRU 16/136/54) using UK aid from the UK Government to support global health research. The views expressed are those of the authors and not necessarily those of the UK Research and Innovation GCRF or the NIHR. The funding sources had no role in the design and content of this paper. Ethical approval was obtained from King's College London Research Ethics Office (reference LRS-19/20–19091).

### ***References***

1. Du R-H, Liang L-R, Yang C-Q, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. *Eur Respir J* 2020;55:2000524.
2. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* 2020;395:497–506.
3. Guan W-j, Ni Z-y, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *New Engl J Med* 2020;382:1708–1720.
4. Sawaya T, Ballouz T, Zaraket H, Rizk N. Coronavirus disease (COVID-19) in the Middle East: a call for a Unified response. *Front Public Health* 2020;8:209.
5. Murthy S, Leligdowicz A, Adhikari NK. Intensive care unit capacity in low-income countries: a systematic review. *PLoS One* 2015;10:e0116949.
6. Reid EA, Kovalerchik O, Jubanyik K, Brown S, Hersey D, Grant L. Is palliative care cost-effective in low-income and middle-income countries? A mixed-methods systematic review. *BMJ Support Palliat Care* 2019;9:120–129.
7. Potts M, Cartmell KB, Nemeth L, Bhattacharjee G, Qanungo S. A systematic review of palliative care Intervention outcomes and outcome measures in low-Resource countries. *J pain symptom Manag* 2018;55:1382–1397.e7.
8. Radbruch L, Knaut FM, de Lima L, de Joncheere C, Bhadelia A. The key role of palliative care in response to the COVID-19 tsunami of suffering. *The Lancet* 2020;395:1467–1469.
9. WHO. Clinical management of COVID-19: interim guidance. World Health Organization, 2020.
10. WHO. 73rd session of the World Health Assembly. COVID-19 response. Resolution 1.7. World Health Organization, 2020.
11. Lovell N, Maddocks M, Etkind SN, et al. Characteristics, symptom management, and outcomes of 101 patients with COVID-19 referred for hospital palliative care. *J Pain Symptom Manage* 2020;60:e77–e81.



12. Arya A, Buchman S, Gagnon B, Downar J. Pandemic palliative care: beyond ventilators and saving lives. *Cmaj* 2020;192:E400–E404.
13. Dubey S, Biswas P, Ghosh R, et al. Psychosocial impact of COVID-19. *Diabetes Metab Syndr* 2020;14:779–788.
14. WHO. Joint external evaluation tool: International health regulations (2005). World Health Organization, 2016.
15. International Health Regulations (2005), Third edition 2016.
16. Etkind SN, Bone AE, Lovell N, et al. The role and response of palliative care and hospice services in epidemics and pandemics: a rapid review to inform practice during the COVID-19 pandemic. *J Pain Symptom Manage* 2020;60:e31–e40.
17. Nouvet E, Sivaram M, Bezanson K, et al. Palliative care in humanitarian crises: a review of the literature. *J Int Humanitarian Action* 2018;3:5.
18. The Lancet. Palliative care and the COVID-19 pandemic. *Lancet* 2020;395:1168.
19. Abubakar A, Elkholy A, Barakat A, et al. Pandemic influenza preparedness (PIP) framework: progress challenges in improving influenza preparedness response capacities in the Eastern Mediterranean Region, 2014–2017. *J Infect Public Health* 2020;13:446–450.
20. Azziz-Baumgartner E, Smith N, González-Alvarez R, et al. National pandemic influenza preparedness planning. *Influenza Other Respir Viruses* 2009;3:189–196.
21. Dhama K, Malik YS, Malik SVS, Singh RK. Ebola from emergence to epidemic: the virus and the disease, global preparedness and perspectives. *The J Infect Developing Countries* 2015;9:441–455.
22. Fineberg HV. Pandemic preparedness and response—lessons from the H1N1 influenza of 2009. *New Engl J Med* 2014;370:1335–1342.
23. Jacobsen KH, Aguirre AA, Bailey CL, et al. Lessons from the Ebola outbreak: Action items for emerging infectious disease preparedness and response. *EcoHealth* 2016;13:200–212.
24. Rajakaruna SJ, Liu WB, Ding YB, Cao GW. Strategy and technology to prevent hospital-acquired infections: lessons from SARS, Ebola, and MERS in Asia and west Africa. *Mil Med Res* 2017;4:32.
25. WHO. IHR monitoring and evaluation framework. International Health Regulations (2005). World Health Organization, 2018.
26. WHO-EMRO. Coronavirus disease 2019 (COVID-19) strategic preparedness and response plan: Accelerating readiness in the Eastern Mediterranean Region. World Health Organ Reg Off East Mediterr, 2020.
27. Nebehay S. Coronavirus could cause upheaval across Middle East - Red Cross. Reuters, 2020.
28. Fadhil I, Lyons G, Payne S. Barriers to, and opportunities for, palliative care development in the Eastern Mediterranean Region. *The Lancet Oncol* 2017;18:e176–e184.
29. Clark D, Baur N, Clelland D, et al. Mapping Levels of palliative care development in 198 countries: the Situation in 2017. *J Pain Symptom Manage* 2020;59:794–807.e4.
30. Asfahan S, Chawla G, Dutt N. Ramadan and COVID-19: a challenge amongst challenges. *Turk Thorac J* 2020;21:285–286.
31. Hosseini Zijoud SR, Jalali Farahani A. Ramadan Coincides with the COVID-19 pandemic: what should Be done?. *Disaster Med Public Health Prep*, 2020:1–2.
32. Javanmard SH, Otraj Z. Ramadan Fasting and risk of covid-19. *Int J Prev Med* 2020;11:60.
33. Waqar S, Ghouri N. Managing Ramadan queries in COVID-19. *BJGP Open* 2020;4:bjgpopen20X101097.
34. Abdullah R, Guo P, Harding R. Preferences and experiences of Muslim patients and their families in Muslim-majority countries for end-of-life care: a systematic review and thematic analysis. *J Pain Symptom Manage* 2020;60:1223–1238.e4.
35. Eysenbach G. Improving the quality of Web surveys: the Checklist for reporting results of internet E-surveys (CHERRIES). *J Med Internet Res* 2004;6:e34.
36. Regmi PR, Waithaka E, Paudyal A, Simkhada P, van Teijlingen E. Guide to the design and application of online questionnaire surveys. *Nepal J Epidemiol* 2016;6:640–644.
37. Arias-Casais N, Garralda E, Rhee J. EAPC Atlas of palliative care in Europe. Romania 2019;122:0–6.
38. Osman H, Rihan A, Garralda E, et al. Atlas of Palliative Care in the eastern Mediterranean region 2017.
39. Rhee JY, Garralda E, Namisango E, et al. The African palliative care association (APCA) Atlas of palliative care development in Africa: a comparative analysis. *The Lancet Glob Health* 2018;6:S21.
40. International Association for Hospice and Palliative Care. Global Directory of Palliative Care Institutions and Organizations. updated 13 April 2020. Available from [www.hospicecare.com/global-directory-of-providers-organizations/](http://www.hospicecare.com/global-directory-of-providers-organizations/).
41. Boufkhed S, Namisango E, Luyirika E, et al. Preparedness of African palliative care services to respond to the COVID-19 pandemic: a rapid assessment. *J Pain Symptom Manage* 2020;60:e10–e26.
42. Costantini M, Sleeman KE, Peruselli C, Higginson IJ. Response and role of palliative care during the COVID-19 pandemic: a national telephone survey of hospices in Italy. *Palliat Med* 2020;34:889–895.
43. Reperant LA, Osterhaus A. AIDS, Avian flu, SARS, MERS, Ebola, Zika... What next? *Vaccine* 2017;35:4470–4474.
44. WHO. Clinical Management of Patients with Viral Haemorrhagic Fever: A Pocket Guide for Front-line Health Workers. Interim Emergency Guidance for Country Adaptation. World Health Organization, 2016.
45. Palagyi A, Marais BJ, Abimbola S, Topp SM, McBryde ES, Negin J. Health system preparedness for emerging infectious diseases: a synthesis of the literature. *Glob Public Health* 2019;14:1847–1868.
46. Ortu G, Mounier-Jack S, Coker R. Pandemic influenza preparedness in Africa is a profound challenge for an already distressed region: analysis of national preparedness plans. *Health Policy Plan* 2008;23:161–169.
47. Sambala EZ, Kanyenda T, Iwu CJ, Iwu CD, Jaca A, Wiysonge CS. Pandemic influenza preparedness in the WHO African region: are we ready yet? *BMC Infect Dis* 2018;18:567.

48. WHO-EMRO. Implementation of the pandemic influenza preparedness (PIP) framework in the Eastern Mediterranean region. Cairo: World Health Organ Reg Off East Mediterr Region, 2017.
49. O’Cathain A, Thomas KJ. "Any other comments?" Open questions on questionnaires - a bane or a bonus to research? *BMC Med Res Methodol* 2004;4:25.
50. Al-Shorbaji N, Househ M, Taweel A, et al. Middle East and North African health Informatics association (MENA-HIA): Building sustainable Collaboration. *Yearb Med Inform* 2018;27:286–291.
51. Moghaddasi H, Mohammadpour A, Bouraghi H, Azizi A, Mazaherilaghab H. Hospital Information Systems: the status and approaches in selected countries of the Middle East. *Electron Physician* 2018;10:6829–6835.
52. Ross J, Stevenson F, Lau R, Murray E. Factors that influence the implementation of e-health: a systematic review of systematic reviews (an update). *Implementation Sci* 2016; 11:146.
53. Calton B, Abedini N, Fratkin M. Telemedicine in the time of coronavirus. *J Pain Symptom Manage* 2020;60: e12–e14.
54. Kirkpatrick JN, Hull SC, Fedson S, Mullen B, Goodlin SJ. Scarce-resource allocation and patient triage during the COVID-19 pandemic: JACC review Topic of the Week. *J Am Coll Cardiol* 2020;76:85–92.
55. Shamieh O, Richardson K, Abdel-Razeq H, Harding R, Sullivan R, Mansour A. COVID-19 - impact on DNR Orders in the Largest cancer center in Jordan. *J pain symptom Manag* 2020;60:e87–e89.
56. Fins JJ. Resuscitating patient rights during the pandemic: COVID-19 and the risk of Resurgent Paternalism. *Cambridge Q Healthc Ethics* 2020:1–15.
57. Fritz Z, Huxtable R, Ives J, Paton A, Slowther AM, Wilkinson D. Ethical road map through the covid-19 pandemic. *Bmj* 2020;369:m2033.

## Appendix I

### Questionnaire in English

Note: Question with an asterisk were compulsory.

COVID-19 preparedness in hospices and palliative care services in the Middle-East and North Africa.

#### ABOUT YOU AND YOUR SERVICE

**\*In which country is your service in?**

- |                                      |                                    |   |
|--------------------------------------|------------------------------------|---|
| <input type="checkbox"/> Afghanistan | <input type="checkbox"/> Kuwait    | <input type="checkbox"/> Qatar                |
| <input type="checkbox"/> Algeria     | <input type="checkbox"/> Lebanon   | <input type="checkbox"/> Saudi Arabia         |
| <input type="checkbox"/> Bahrain     | <input type="checkbox"/> Libya     | <input type="checkbox"/> Syria                |
| <input type="checkbox"/> Egypt       | <input type="checkbox"/> Morocco   | <input type="checkbox"/> Tunisia              |
| <input type="checkbox"/> Iran        | <input type="checkbox"/> Oman      | <input type="checkbox"/> Turkey               |
| <input type="checkbox"/> Iraq        | <input type="checkbox"/> Palestine | <input type="checkbox"/> United Arab Emirates |
| <input type="checkbox"/> Jordan      | <input type="checkbox"/> Pakistan  | <input type="checkbox"/> Yemen                |

**Characteristics of your hospice/palliative care service:**

- Approximate number of patients seen per year:
- \*Type of hospice and/or service: *(tick all that applies)*
  - Private
  - Non-profit charity
  - Government or public.
  - Within a hospital
  - Within community
  - Inpatient hospice
  - Outpatient hospice
  - Other, please specify:
- Do you have beds:  Yes  No

**IF YES**, Number of beds:

**\*What is your current role?** *(tick all that applies)*

- Doctor or medical officer.
- Nurse.
- Psychosocial professional.
- Manager or responsible of the hospice or palliative care service.
- Other, please specify:

#### CURRENT COVID-19 SITUATION IN YOUR SERVICE

1. **\*Did you have a suspected or confirmed cases of COVID-19 in your service or in the hospital you are based in?**

- yes, confirmed cases  yes, probable cases  yes, suspect cases  no (If no, go to Q.2)

IF YES:

- **Who was positive?** (tick all that applies)
  - patients.
  - relatives.
  - physicians.
  - nurses.
  - volunteers.
  - other (e.g. administrative or cleaning staff), please specify:
- **How many cases did you have (specify numbers for suspected, probable, confirmed)?**
- **Where were the cases identified?**
  - your service or hospice  another service of the hospital you are based in.
- **How were they identified?** (e.g. who informed you, which communication means (phone, email, etc.)
- **What was done?** (e.g. reporting, referral, containment measures, protection of and communication with staff and users, etc.)
- **What were the consequences?** (e.g. for your service, yourself, your interaction with the community, etc.)

#### WRITTEN PROCEDURES (OR GUIDANCE)

2.1 **\*Do you have a case definition for confirmed, probable and suspected COVID-19 cases?**

- yes  no  do not know/not sure.

**2.2 Do you have a written procedure for “what to do” if you have a confirmed, probable and/or suspected COVID-19 case in your service among:**

	Procedure Specific to COVID-19	Procedure for Infectious Diseases in General or to Another Specific Highly Infectious Disease (e.g. Influenza, Ebola, Tuberculosis, etc.) Please specify for which disease(s):
- patients	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know
- relatives and visitors	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know
- healthcare professional staff member	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know
- volunteers and medical staff	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know
- staff and volunteers going in the community	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know
- Other, please specify:	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> don't know

**2.3 Do you have a procedure to support healthcare providers to manage stress?  yes  no  don't know**

(optional) Please specify or comment:

(optional) Additional thoughts on policies and protocols?

*MEASURES IN PLACE (TO AVOID CONTAGION)*

**3.1 Did you modify any of the following policies or procedures as a measure to avoid contagion?**

- Policy for visitors/relatives (number of visitors, hours etc.)	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> not sure <input type="checkbox"/> N/A
- Policy for operator protection (personal protective equipment)	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> not sure <input type="checkbox"/> N/A
- Policy for patients' admission to the hospice	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> not sure <input type="checkbox"/> N/A
- Volunteer support policy	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> not sure <input type="checkbox"/> N/A
- Policy on how to handle dead patients	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> not sure <input type="checkbox"/> N/A
- 'Do Not Resuscitate' (DNR) policy	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> not sure <input type="checkbox"/> N/A
- Policy regarding care of the relatives after the patient's death	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> not sure <input type="checkbox"/> N/A
- Other policy modified, please specify:	<input type="checkbox"/> yes <input type="checkbox"/> no <input type="checkbox"/> not sure <input type="checkbox"/> N/A

Please tick N/A if you do not have the corresponding policy or procedure in place.

**IF YOU ANSWERED YES TO ANY OF THE ABOVE:**

Did you change the policies following the instructions from health management or regional authorities, or did your hospice take them spontaneously?

following the instructions  spontaneously  both.

**3.2 Do you have in place any of the following measures to protect staff and patients:**

- **Hand washing facility for all at points of entry** (soap and running water or hand sanitizers with 60% alcohol):  
 Yes, we put additional ones  We already had them in place before COVID-19  No, we do not have such facility.
- **Personal Protection Equipment (PPE) for:**
  - o **palliative care staff:**  Yes, we put additional ones  Not more than usual  No, we do not have PPE
  - o **cleaning staff:**  Yes, we put additional ones  Not more than usual  No, we do not have PPE
- **Isolation room identified in case of infectious conditions, like COVID-19:**  Yes  No  N/A (outpatient service only)
- **Recommendations if you or someone in your household becomes ill with COVID-19 symptoms**  
 yes  no  don't know.

*IF YOU ANSWERED YES TO ANY OF THE ABOVE:*

Did you put the measure in place following the instructions from health management or regional authorities, or did your hospice take them spontaneously?

following the instructions  spontaneously  both.

**3.3 Have all healthcare providers been trained in handling highly infectious conditions such as COVID-19?**

Yes, trained before COVID-19 pandemic  Yes, trained because of COVID-19 pandemic  Not trained.

**3.4 Do you know how to dispose of highly infectious waste?**

- in the hospice or service  Yes  No  N/A (outpatient service only)
- in the community  Yes  No  N/A (inpatient/hospice service only)

### 3.5. Was the cleaning staff included in information sharing and training regarding managing COVID-19?

(e.g. adapting practice in case of COVID-19 suspected)  Yes  No  Don't know/Not sure.  
(optional) Additional thoughts on measures in place to avoid contagion:

## COMMUNICATION AND COORDINATION

### 4.1 How would you be informed if there is a confirmed or suspected case in the hospice or in the locality?

- Who or which institution will inform your hospice or service?
- Who will be informed in your hospice or service (position or job title)?
- Communication system(s) that will be used (tick all that applies)
  - Mobile phone available 24/7.
  - Telephone (in the service)
  - Email.
  - WhatsApp/Viber group.
  - Other, please specify:

### 4.3 Is there a focal point person identified in the hospice or service responsible for collecting and sharing up-to-date information (about health recommendations, cases, protocols to use): yes no not sure

Please specify (job title/position):

### 4.4 What communication means are in place to share COVID-19 or other urgent information with ...

(tick all that applies)

- the staff?  Text message  WhatsApp/Viber  Phone call  Email  None  Other, specify
- patients?  Text message  WhatsApp/Viber  Phone call  Email  None  Other, specify
- relatives, visitors?  Text message  WhatsApp/Viber  Phone call  Email  None  Other, specify

### 4.5 Do you have an up-to-date contact list of ...

- |  |  |  |                             |                                 |
|--|--|--|-----------------------------|---------------------------------|
| -all staff working in or for the hospice or service (medical, administrative, cleaning staff, etc.)? | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |
| -all patients that attended or have attended the hospice or service?                                 | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |
| -all relatives that visited or have visited the hospice or service?                                  | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |
| -patients visited in the community?  | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |

### 4.6 Do you have a system collecting information about ...

- |                                     |  |  |                             |                                 |
|-------------------------------------|--|--|-----------------------------|---------------------------------|
| -Patients' symptoms?                | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |
| -Patients' outcomes?                | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |
| -Treatment given?                   | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |
| -Dates of patients' visits or stay? | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |
| -Dates of relatives' visits?        | <input type="checkbox"/> Yes, a paper-based registry | <input type="checkbox"/> Yes, an electronic record | <input type="checkbox"/> No | <input type="checkbox"/> Other: |

(optional) Additional thoughts on communication and coordination?

## 5. RESOURCES

### 5.1 Do you have concerns about access to ...

	In Your Hospice or Service?	In the Surrounding Community?
-running water?	<input type="checkbox"/> yes <input type="checkbox"/> no	<input type="checkbox"/> yes <input type="checkbox"/> no
-soap?	<input type="checkbox"/> yes <input type="checkbox"/> no	<input type="checkbox"/> yes <input type="checkbox"/> no
-hand sanitizers (with at least 60% alcohol)?	<input type="checkbox"/> yes <input type="checkbox"/> no	<input type="checkbox"/> yes <input type="checkbox"/> no
-electricity?	<input type="checkbox"/> yes <input type="checkbox"/> no	<input type="checkbox"/> yes <input type="checkbox"/> no
-thermometers (contactless, <i>Thermoflash</i> -type)?	<input type="checkbox"/> yes <input type="checkbox"/> no	<input type="checkbox"/> yes <input type="checkbox"/> no
-accessing disinfectant products to continue providing care safely?	<input type="checkbox"/> yes <input type="checkbox"/> no	<input type="checkbox"/> yes <input type="checkbox"/> no
-other, please specify:	<input type="checkbox"/> yes <input type="checkbox"/> no	<input type="checkbox"/> yes <input type="checkbox"/> no

### 5.2 Do you have adequate material and facilities to dispose of highly infectious waste ...

- in the hospice?  yes  no  don't know/not sure
- in the community?  yes  no  don't know/not sure

### 5.3 Do you have an up-to-date inventory of ...

- protection material available for staff, patient and visitors (*hygiene and sanitation materials, protection material like masks, etc.*)?  yes  no  not sure
- medicines and other medical supplies available to care for the patients?  yes  no  not sure

**5.4 Do you have the capacity to use technology instead of face-to-face appointment to provide some care remotely?**  yes  no

**IF YES**, please specify:

- which technology? (tick all that applies):  Phone call  Video call  Other, please specify:
- which service can be provided remotely? (e.g. *psychological support, spiritual care, grief and bereavement, managing the end of life phase, etc.*)
- what has worked well when using virtual technologies?
- what was difficult when using technologies?

**IF NO**, please specify:

- what are your limitations to use technology?
- what would facilitate your use of technology?

**5.5 In case of emergency, lockdown or quarantine, do you know how your hospice would access to:** (e.g. Local/national authorities' stocks, private supplier, transportation, etc.)

- food (*for inpatient services only*)?  yes  no
  - medicines and other medical supply?  yes  no
  - additional staff (e.g. *if staff self-isolates or becomes ill*)?  yes  no
- (optional) Please specify:

**5.6 Do you have education material about COVID-19 available?**  yes  no

**IF YES**, are there posters displayed where staff, patients and visitors can see them?  yes  no  
are they also available for the surrounding community?  yes  no

Please specify:

- Which education material do you have?
- How you did you get the education material?

(optional) Additional thoughts on resources:

## 6. EFFECTS ON STAFF

**6.1 Did you observe that some staff suddenly did not come to work without justification**

(i.e. *more than usual*)?  yes  no  not sure.

**6.2 In your opinion, how anxious are your staff about the need to care for their children who may not be at school? From 1 to 10 (1-not at all anxious; 10-extremely anxious)**

1 (Not at all)  2  3  4  5  6  7  8  9  10 (extremely)

**6.3 In your opinion, how anxious are your staff about the need to care for their own relatives?**

From 1 to 10 (1-not at all anxious; 10-extremely anxious)

1 (Not at all)  2  3  4  5  6  7  8  9  10 (extremely)

**6.4 In your opinion, how anxious are your staff about getting infected themselves?**

From 1 to 10 (1 – not at all anxious; 10-extremely anxious)

1 (Not at all)  2  3  4  5  6  7  8  9  10 (extremely)

**6.5 How worried are you regarding potential issues for your interaction with the community if your hospice or service is known to manage a potential COVID-19 case?**

From 1 to 10 (1-not at all worried; 10-extremely worried)

1 (Not at all)  2  3  4  5  6  7  8  9  10 (extremely)

(optional) Additional thoughts on other potential effects of the COVID-19 on you and your staff:

### 7. PERCEPTION OF THE RISK

In the coming week ....

**7.1 How much do you think hospice/palliative care staff are at risk of being infected by COVID-19?**

From 0–10 (0 no risk - 10 maximum risk you can imagine)

1 (none)  2  3  4  5  6  7  8  9  10 (maximum)

**7.2 How much do you think the hospice/palliative care service is at risk of closing because of an infection in the hospice or service? From 0–10 (0 no risk - 10 maximum risk you can imagine)**

1 (none)  2  3  4  5  6  7  8  9  10 (maximum)

**7.3 Do you have any security concerns for yourself or your staff?**  yes  no

IF YES, please specify ...

(optional) Additional thoughts on other potential effects of the COVID-19 on your staff:

### 8. PREPARING TO OFFER SUPPORT

**8.1 Do you have palliative care protocols for symptom management and psychological support that could be shared with non-specialist staff and/or COVID-19 response teams in other healthcare facilities?**  yes  no

**IF YES, do you have the capacity to train non-specialist in using these protocols?**  yes  no

Optional: what are your limitations to share your expertise?

Optional: what could facilitate the sharing of your expertise?

**8.2 In case of outbreak COVID-19 or another highly infectious disease, do you have plans to redeploy the following outside of the inpatient settings?**

-Healthcare providers	<input type="checkbox"/> yes	<input type="checkbox"/> no	<input type="checkbox"/> don't know	<input type="checkbox"/> N/A
-Volunteers	<input type="checkbox"/> yes	<input type="checkbox"/> no	<input type="checkbox"/> don't know	<input type="checkbox"/> N/A
-Resources (material and supplies)	<input type="checkbox"/> yes	<input type="checkbox"/> no	<input type="checkbox"/> don't know	<input type="checkbox"/> N/A

**IF YES to any of the above:**

**Please specify in which settings** they be re-deployed (e.g. community settings, another service, etc.)?

**8.3 Do you have plans to support other healthcare services in the triage of patients in case of COVID-19 outbreak?**  yes  no

(optional) Comment:

### ADDITIONAL COMMENTS

What do you foresee will be the biggest challenges for COVID-19 in your service over the next 1–2 months?

What would help you most to overcome these?

Do you think there are relevant information we have omitted to ask you?

What are your biggest worries or concerns?

Feel free to share any additional thought or comment:

### OPTIONAL:

**Would you like to receive the results of this survey via e-mail?**  yes  no

**Would you like to be contacted in the future about opportunities about research on or advocacy for palliative care?**  yes  no

**If yes to any of the above, please share your contact details (name, organization, email):**

Please note: We will separate this information from your responses to the questions, and only selected team members at King's College London (KCL) will access it. Please note that if you agree to share your contact details, your reply will no longer be anonymous to the KCL research team. Your answers will be treated confidentially, and your data will be held securely.

*FINAL INFORMATION***You can find information about COVID-19 and Palliative Care in the following resources:**

- Cicely Saunders Institute for Palliative Care and Rehabilitation, King's College London [www.kcl.ac.uk/cicelysaunders/resources/links](http://www.kcl.ac.uk/cicelysaunders/resources/links)
- Worldwide Hospice Palliative Care Alliance [www.thewhpc.org/covid-19](http://www.thewhpc.org/covid-19)
- European Association for Palliative Care [www.eapcnet.eu/publications/coronavirus-and-the-palliative-care-response](http://www.eapcnet.eu/publications/coronavirus-and-the-palliative-care-response)

**You can find information about COVID-19 in the following resources:**

- World Health Organization (WHO) [www.who.int/emergencies/diseases/novel-coronavirus-2019](http://www.who.int/emergencies/diseases/novel-coronavirus-2019)
- WHO EMRO [www.emro.who.int/health-topics/corona-virus/information-resources.html](http://www.emro.who.int/health-topics/corona-virus/information-resources.html)
- WHO EURO [www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19](http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19)

**If you have questions or concerns, please contact Sabah Boufkhed: [sabah.boufkhed@kcl.ac.uk](mailto:sabah.boufkhed@kcl.ac.uk).**

Thank you very much for your time and for taking part in the survey.



## Appendix II. Questionnaire in Arabic

جاهزية دور الرعاية وخدمات العناية التلطيفية لفيروس كورونا المستجد كوفيد-19 في الشرق الأوسط وشمال إفريقيا

### أسئلة عنك وعن الوحدة التي تعمل بها

\*في أية دولة تقع الوحدة التي تعمل بها؟

قطر	الكويت	أفغانستان
السعودية	لبنان	الجزائر
سوريا	ليبيا	البحرين
تونس	المغرب	مصر
تركيا	عمان	إيران
الإمارات	فلسطين	العراق
اليمن	الباكستان	الأردن

مواصفات دار الرعاية/وحدة العناية التلطيفية:

عدد المرضى الذين يتم معالجتهم سنوياً: ...

\*نوع دار الرعاية أو وحدة العناية التلطيفية: (قم باختيار جميع الخيارات المنطبقة على إجابتك):

خاص

خيري غير ربحي

حكومي

جزء من مستشفى

مجتمعي

دار رعاية للمرضى المنومين

عيادات دار رعاية

أخرى، قم بذكرها:

نعم/لا

هل هنالك أسرة في الوحدة التي تعمل بها؟

في حال كانت الإجابة نعم، لطفاً قم بكتابة عدد الأسرة:

\* ما هو دورك الحالي؟

(قم باختيار جميع الخيارات المنطبقة على إجابتك)

طبيب أو مسؤول طبي/ممرض/مدير أو مسؤول عن دار الرعاية أو وحدة العناية التنطيفية / الأخصائي النفسي والاجتماعي/أخرى، قم بذكرها

### 1. الوضع الوبائي لفايروس كورونا المستجد كوفيد-19 في الوحدة التي تعمل بها

1. \* هل لديك حالة مؤكدة أو مشكوك بإصابتها بفايروس كورونا المستجد كوفيد-19 في الوحدة أو دار

الرعاية التي تعمل بها؟

نعم، حالة مؤكدة

نعم، حالة محتملة

نعم، حالة مشكوك بإصابتها

لا

لست متأكد، لطفاً قم بذكرها...

إذا كانت الإجابة لا، انتقل إلى السؤال الثاني.

إذا كانت الإجابة نعم (حالة مؤكدة، حالة محتملة ، حالة مشكوك بإصابتها):

من هم المشخصون بفايروس كورونا المستجد كوفيد-19؟

(قم باختيار جميع الخيارات المنطبقة على إجابتك)

مرضى

أقرباء المرضى

أطباء

ممرضون

متطوعون

أخرى (كالإداريين أو طواقم التنظيف)، قم بذكرها

أين تم تشخيص الحالات؟

في الوحدة أو دار الرعاية التي تعمل بها

وحدة أخرى في ذات المستشفى الذي تعمل به

كم حالة تواجدت لديك؟

(قم بذكرها بالتفصيل من حيث المشكوك بإصابتهم أو المحتمل إصابتهم أو المؤكد إصابتهم).

كيف تم التقصي عنهم؟

(مثال: من قام بتبليغك؟ ما هي طريقة التبليغ؟ عبر الهاتف أم عبر البريد الإلكتروني؟ إلخ).

ماذا كان الإجراء المتبع؟

(مثال: التقرير، التحويل، إجراءات الاحتواء، حماية الطاقم والتواصل معهم، إلخ)

ماذا كانت التبعات؟

(مثال: التبعات على وحدتك، عليك شخصياً، على تعاملك مع المجتمع، إلخ)

## 2. الإجراءات أو الإرشادات الخطية

2.1 \* هل لديكم تعريف للحالات المؤكد إصابتها والمحتمل إصابتها والمشكوك إصابتها بفايروس كورونا المستجد كوفيد-

؟19

نعم

لا

لا أعرف / لست متأكد

2.2 هل لديكم إجراءات خطية للتعامل مع الحالات المؤكد إصابتها والمشكوك بإصابتها والمحتمل إصابتها في الوحدة التي

تعمل بها لكل من

إجراء لكوفيد-19 على وجه الخصوص	إجراء للأمراض المعدية بشكل عام أو لأمراض معينة معدية بشكل كبير (كالانفلونزا أو الإيبولا أو السل)*  لطفاً قم بتحديد المرض ...	
المرضى	نعم/لا/لا اعلم	نعم/لا/لا اعلم
الأقارب والزوار	نعم/لا/لا اعلم	نعم/لا/لا اعلم
موظفو الرعاية الصحية	نعم/لا/لا اعلم	نعم/لا/لا اعلم
المتطوعون والطواقم الطبية	نعم/لا/لا اعلم	نعم/لا/لا اعلم
الموظفون والمتطوعون المخالطون للمجتمع	نعم/لا/لا اعلم	نعم/لا/لا اعلم
أخرى، *قم بذكرها	نعم/لا/لا اعلم	نعم/لا/لا اعلم

2.3 هل لديكم إجراءات لدعم موظفي الرعاية الصحية للتعامل مع التوتر والقلق؟ نعم/لا/لا أعلم

(اختياري) لطفاً أنكرها وأكتب تعليقاً

(اختياري) قم بذكر أية أفكار أو اقتراحات فيما يتعلق بالسياسات والبروتوكولات

### 3. الإجراءات المتبعة لتجنب التفشي

3.1 هل قمت بتعديل أي من السياسات أو الإجراءات التالية بهدف تجنب التفشي؟

لطفاً قم بالإشارة على (غير متوفر) إذا لم تكن السياسة أو الإجراء متبعاً لديكم

سياسات الزوار والأقارب (كعدد الزوار وساعات الزيارة)	نعم/لا/لست متأكد/غير متوفر
سياسات حماية العاملين (معدات الوقاية الشخصية)	نعم/لا/لست متأكد/غير متوفر
سياسات إدخال المرضى إلى دار الرعاية	نعم/لا/لست متأكد/غير متوفر
سياسات التطوع	نعم/لا/لست متأكد/غير متوفر
سياسات التعامل مع المرضى المتوفين	نعم/لا/لست متأكد/غير متوفر
سياسات عدم الانعاش	نعم/لا/لست متأكد/غير متوفر
سياسات العناية بأقارب المتوفى بعد وفاته	نعم/لا/لست متأكد/غير متوفر
سياسات أخرى، لطفاً قم بذكرها...	

في حال كانت إجابتك "نعم" لأي من الأسئلة أعلاه:

هل قمت بتغيير السياسات تبعاً لتعليمات الإدارة الصحية أو السلطات المحلية، أم أن دار الرعاية التي تعمل بها قامت بتعديلها تلقائياً؟

تبعاً للتعليمات/بشكل تلقائي/لكلا السببين

3.2 هل يتم اتباع أي من الإجراءات التالية لحماية الطاقم والمرضى:

-محطات تغسيل اليدين في كافة نقاط الدخول (الصابون والماء أو المعقمات اليدوية بنسبة كحول 60%:

نعم، قمنا بإضافة محطات أخرى/كانت متوفرة قبل وباء كوفيد-19/لا، ليست لدينا أية محطات كالمذكورة أعلاه

-معدات الوقاية الشخصية لكل من:

\*طاقم العناية التلطيفية: نعم، قمنا بإضافة معدات إضافية/لا شيء أكثر من المعتاد/لا، ليست لدينا

معدات وقاية شخصية

\*طاقم التنظيف: نعم، قمنا بإضافة معدات إضافية/لا شيء أكثر من المعتاد/لا، ليست لدينا

معدات وقاية شخصية

-غرف العزل للحالات المعدية ككوفيد-19: نعم/لا/غير متوفر (للعيادات الخارجية)

-توصيات لك أو لأحد أهل بيتك في حال إصابته بأعراض كوفيد-19: نعم، لطفاً قم بذكرها/لا/لا أعلم

إذا كانت إجابتك "نعم" لأي من المذكور أعلاه:

هل قمت بتنفيذ الإجراءات تبعاً لتعليمات الإدارة الصحية أو السلطات المحلية، أم أن دار الرعاية التي تعمل بها قامت

بتعديلها تلقائياً؟  
تبعاً للتعليمات/بشكل تلقائي/لكلا السببين

3.3 هل تم تدريب جميع موظفي الرعاية الصحية على كيفية التعامل مع الحالات شديدة العدوى ككوفيد-19؟

نعم، تم تدريبهم قبل وباء كوفيد 19 /نعم، تم تدريبهم تبعاً لوباء كوفيد-19/لم يتم تدريبهم

### 3.4 هل أنت على علم بكيفية التخلص من النفايات عالية العدوى؟

-في دار الرعاية أو الوحدة التي تعمل بها: نعم/لا/غير متوفر

-في المجتمع: نعم/لا/غير متوفر

### 3.5 هل تم تدريب طواقم التنظيف وإطلاعهم على المعلومات المتعلقة بالتعامل مع وباء كوفيد-19؟

(التكيف مع التعامل مع حالات كوفيد-19 المشكوك بها)

نعم/لا/غير متوفر / لا أعرف / لست متأكدًا

(اختياري) هل لديك أية أفكار أو اقتراحات بخصوص الإجراءات المتبعة لتجنب التفشي؟

## 4. التواصل والتنسيق

### 4.1 كيف يتم تبليغكم بوجود حالات مؤكدة أو مشكوك بأمرها في دار الرعاية أو في الجوار؟

-ما هي الجهة أو المؤسسة التي تقوم بتبليغ دار الرعاية أو الوحدة التي تعمل بها؟

-من هو المبلِّغ بالأمر في دار الرعاية أو الوحدة التي تعمل بها؟ (مسماه الوظيفي أو منصبه)

-نظم التواصل المتبعة (قم باختيار جميع الخيارات المنطبقة على إجابتك)

( ) هاتف نقال متوفر على مدار الساعة

( ) هاتف متوفر في الوحدة

( ) بريد الكتروني

( ) مجموعة على تطبيق الواتساب أو الفاير

( ) أخرى، لطفًا قم بذكرها...

4.3 هل هنالك موظف ارتباط في دار الرعاية أو الوحدة الخاصة بك معني بجمع ومشاركة أحدث المعلومات؟

(تلك المتعلقة بالتوصيات الصحية أو الحالات أو البروتوكولات المتبعة) نعم/لا/لست متأكدًا

لطفًا قم بذكر منصبه أو مسماه الوظيفي:

4.4 ما هي وسائل التواصل المتبعة لمشاركة المعلومات المتعلقة بكوفيد-19 أو أية معلومات طارئة مع كل من:

(قم باختيار جميع الخيارات المنطبقة على إجابتك)

**-الطاقم:** رسالة نصية/واتساب/فايبر/مكالمة

هاتفية/بريد الكتروني/لا يوجد وسائل/أخرى، قم بذكرها أدناه

**-المرضى:** رسالة نصية/واتساب/فايبر/مكالمة هاتفية/بريد الكتروني/لا يوجد وسائل/أخرى، قم بذكرها أدناه

**-الأقارب والزوار:** رسالة نصية/واتساب/فايبر/مكالمة هاتفية/بريد الكتروني/لا يوجد وسائل/أخرى، قم بذكرها

أدناه

4.5 هل لديك قائمة جهات اتصال محدثة لكل من:

**-الطاقم، سواء العاملين في دار الرعاية أو الوحدة أو العاملين لصالحها (الطواقم الطبية أو الإدارية أو طواقم التنظيف**

إلخ):

نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى...

**-المرضى الذين حضروا إلى دار الرعاية أو الوحدة:**

نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى

**-الأقارب الذين قاموا بزيارة دار الرعاية أو الوحدة:**

نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى

**-المرضى الذين تمت زيارتهم في المجتمع:**



نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى

4.6 هل لديك نظام لجمع المعلومات لكل من:

-أعراض المرضى:

نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى

-نتائج المرضى:

نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى

-العلاجات المعطاة:

نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى

-تواريخ مكوث المرضى:

نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى

-تواريخ زيارة الأقارب:

نعم، يوجد سجل ورقي/نعم، يوجد سجل الكتروني/لا/أخرى

(اختياري) هل لديك أية أفكار أو اقتراحات متعلقة بالتواصل والتنسيق؟

## 5. الموارد

5.1 هل لديك قلق أو شكوك فيما يتعلق بتوفر...

في المجتمع المحيط	في دار الرعاية أو الوحدة	
نعم/لا	نعم/لا	الماء
نعم/لا	نعم/لا	الصابون
نعم/لا	نعم/لا	معقمات الأيدي (بنسبة كحول لا تقل عن 60%)
نعم/لا	نعم/لا	الكهرباء
نعم/لا	نعم/لا	موازين الحرارة (بتقنية القياس عن بعد)
نعم/لا	نعم/لا	المواد المعقمة لضمان توفير الرعاية الصحية بشكل آمن
نعم/لا	نعم/لا	أخرى، لطفاً قم بذكرها

5.2 هل لديك مواد و منشآت كافية للتخلص من النفايات شديدة العدوى؟

-في دار الرعاية: نعم/لا/لا أعلم، لست متأكد

-في المجتمع: نعم/لا/لا أعلم، لست متأكد

5.3 هل لديك خزانة حديثة من كل من:

-مواد الحماية المتوفرة لكل من الطاقم والمرضى والزوار (مواد النظافة والتعقيم أو مواد الوقاية كالكمامات

إلخ): نعم/لا/لست متأكد

-الأدوية والمعدات الطبية الأخرى المتوفرة لرعاية المرضى: نعم/لا/لست متأكد

5.4 هل لديك الإمكانيات لاستخدام التكنولوجيا كبديل للمواعيد وجهاً لوجه من باب توفير شكل من أشكال الرعاية عن

بعد؟: نعم/لا

إذا كانت الإجابة نعم، لطفاً قم بالإجابة عن التالي:

- ما هي التقنية المستخدمة؟ (قم باختيار جميع الخيارات المنطبقة على إجابتك): مكالمة هاتفية/مكالمة فيديو/أخرى، لطفاً قم بذكرها
- ما هي الخدمات التي يمكن توفيرها عن بعد؟ (كالدعم النفسي أو الرعاية الروحانية أو الحزن والفجعة أو الرعاية في الأيام الأخيرة إلخ)
- ما هي إيجابيات استخدام التقنيات الافتراضية؟
- ما هي مصاعب استخدام هذه التقنيات؟

إذا كانت الإجابة لا، لطفاً قم بالإجابة عن التالي:

- ما هي المحددات التي تحول دون استخدام هذه التقنيات؟
  - كيف يمكن تسهيل استخدامك لهذه التقنيات؟
- 5.5 في حالات الطوارئ أو الإغلاق العام أو الحجر الصحي، هل تعمل كيف ستتمكن دار الرعاية التي تعمل بها من توفير:

(سواء من خلال مخزونات السلطات المحلية أو الوطنية أو من مزودي القطاع الخاص أو جهات النقل إلخ)

- الأغذية (للمرضى المقيمين في المستشفى فقط)؟ نعم/لا
- الأدوية والمعدات الطبية الأخرى؟ نعم/لا
- طواقم إضافية (في حال قام إصابة بعض أفراد الطاقم أو قيامهم بالعزل الذاتي)؟ نعم/لا

(اختياري) لطفاً قم بالتوضيح

5.6 هل لديك أية مواد تنقيفية فيما يتعلق بكوفيد-19؟ نعم/لا

إذا كانت إجابتك نعم:

- هل يتم عرض هذه المنشورات في مكان مرئي للطاقم والمرضى والزوار؟ نعم/لا
- هل هي متوفرة كذلك في المجتمع المحيط؟ نعم/لا

**لطفاً قم بتوضيح التالي:**

- ما هي طبيعة المواد التثقيفية لديك؟
- كيف حصلت على هذه المواد التثقيفية؟

(اختياري) قم بذكر أية أفكار أو اقتراحات فيما يتعلق بالموارد

**6. التأثير على الطاقم**

6.1 هل لاحظت تغيب بعض أفراد الطاقم بشكل مفاجئ عن العمل دون بيان السبب (بشكل أكثر من المعتاد)؟

نعم/لا/لست متأكد

6.2 برأيك، ما مدى توتر الطاقم لديك فيما يتعلق بحاجتهم لرعاية أطفالهم من غير الذاهبين إلى المدارس؟ (على مقياس

من 1 إلى 10، حيث 1=غير قلق أبداً، 10=قلق بشدة)

1/2/3/4/5/6/7/8/9/10

6.3 برأيك، ما مدى توتر الطاقم لديك فيما يتعلق بحاجتهم لرعاية أقرانهم؟

(على مقياس من 1 إلى 10، حيث 1=غير قلق أبداً، 10=قلق بشدة)

1/2/3/4/5/6/7/8/9/10

6.4 برأيك، ما مدى قلق الطاقم لديك فيما يتعلق بإصابتهم أنفسهم؟

(على مقياس من 1 إلى 10، حيث 1=غير قلق أبداً، 10=قلق بشدة)

1/2/3/4/5/6/7/8/9/10

6.5 ما مدى قلقك بشأن المشاكل المترتبة على تواصلك مع المجتمع المحيط بك في حال كانت دار الرعاية أو الوحدة التي تعمل بها معروفة بتعاملها مع حالات كوفيد-19 محتملة؟

(على مقياس من 1 إلى 10، حيث 1=غير قلق أبداً، 10=قلق بشدة)

1/2/3/4/5/6/7/8/9/10

(اختياري) قم بذكر أية أفكار أو اقتراحات فيما يتعلق بالآثار المحتملة لكوفيد-19 عليك وعلى طاقمك

### 7. إدراك المخاطر

في غضون الأسبوع القادم...

7.1 برأيك، ما مدى فرصة إصابة طاقم دار الرعاية أو وحدة الرعاية التلطيفية بكوفيد-19؟

(على مقياس من 0 إلى 10، حيث 0=ليس هنالك خطورة، 10=أقصى خطورة يمكنك تخيلها)

1/2/3/4/5/6/7/8/9/10

7.2 برأيك، ما احتمال إغلاق دار الرعاية أو وحدة الرعاية التلطيفية التي تعمل بها بسبب وجود حالة مصابة فيها؟

(على مقياس من 0 إلى 10، حيث 0=ليس هنالك خطورة، 10=أقصى خطورة يمكنك تخيلها)

1/2/3/4/5/6/7/8/9/10

7.3 هل لديك أية مخاوف أمنية فيما يتعلق بك أو بالطاقم؟ نعم/لا

إذا كانت إجابتك نعم، لطفاً قم بالتوضيح

(اختياري)

قم بذكر أية أفكار أو اقتراحات فيما يتعلق بأية آثار أخرى محتملة لكوفيد-19 على طاقمك

## 8. الجاهزية لتقديم الدعم

8.1 هل لديك أية بروتوكولات للعناية التلطيفية فيما يتعلق بالسيطرة على الأعراض والدعم النفسي ويمكن مشاركتها مع

غير الاختصاصيين و/أو فرق الاستجابة لكوفيد-19 في مؤسسات الرعاية الصحية الأخرى؟

نعم/لا

إذا كانت الإجابة نعم، هل لديك الإمكانيات لتدريب غير الاختصاصيين على كيفية تطبيق هذه البروتوكولات؟ نعم/لا

(اختياري)

ما هي المحددات التي تحول دون مشاركة هذه الخبرات؟

(اختياري)

كيف يمكن تسهيل مشاركة هذه الخبرات؟

8.2 في حال حدوث جائحة لكوفيد-19 أو أي مرض معدٍ آخر، هل لديك خطط لنشر الطواقم التالية خارج المستشفيات؟

-موفري الرعاية الصحية: نعم/لا/لا أعلم/غير متوفر

-المتطوعين: نعم/لا/لا أعلم/غير متوفر

-الموارد (المواد واللوازم): نعم/لا/لا أعلم/غير متوفر

إذا أُجبت بنعم على أي من المذكور أعلاه، لطفاً قم بتوضيح السياق الذي سيتم نشرها فيه (كالمجتمع أو

وحدات العناية الأخرى إلخ)

8.3 هل لديك أية خطط لدعم وحدات الرعاية الصحية الأخرى في تصنيف الحالات في حال حدوث جائحة لكوفيد-19؟

نعم/لا

(اختياري) قم بذكر أية تعليقات لديك

تعليقات إضافية

برأيك، ما هي أبرز التحديات المستقبلية لكوفيد-19 في وحدتك على مدى الشهر أو الشهرين القادمين؟

ما هي أفضل الطرق لمساعدتك على تجاوز هذه التحديات؟

هل تعتقد أن هنالك أي معلومات مهمة لم نسألك عنها في هذا الاستبيان؟

ما هي أبرز مخاوفك؟

بإمكانك مشاركة أية أفكار أو تعليقات:

اختياري:

هل ترغب في تلقي نتائج هذا الاستطلاع عبر البريد الإلكتروني؟ نعم/لا

هل ترغب في أن يتم الاتصال بك في المستقبل بشأن فرص البحث أو مناصرة الرعاية التلطيفية؟ نعم/لا

إذا كانت الإجابة بنعم على أي مما ورد أعلاه ، فيرجى تزويدنا بتفاصيل الاتصال (الاسم ، المنظمة ، البريد الإلكتروني) :

ملاحظة:

سيتم تفرقة بياناتكم الشخصية من أجوبتكم على الأسئلة و سيسمح فقط لأعضاء معينين من فريق العمل في كلية كينجز في

لندن الاطلاع على هذه البيانات.

يرجى الملاحظة أنه في حال موافقتكم على مشاركة معلومات التواصل الخاصة بك فإن أجوبتك لن تكون مجهولة لفريق

العمل في كلية كينجز في لندن.

### المعلومات النهائية

بإمكانك الوصول إلى معلومات متعلقة بكوفيد-19 والعناية التلطيفية في الموارد التالية:

-معهد سيسيلي سوندرز للعناية التلطيفية وإعادة التأهيل في كلية الملك في لندن:

[www.kcl.ac.uk/cicelysaunders/resources/links](http://www.kcl.ac.uk/cicelysaunders/resources/links)

-اتحاد الرعاية التلطيفية العالمي

[www.thewhpc.org/covid-19](http://www.thewhpc.org/covid-19)

-المؤسسة الأوروبية للرعاية التلطيفية

[www.eapcnet.eu/publications/coronavirus-and-the-palliative-care-response](http://www.eapcnet.eu/publications/coronavirus-and-the-palliative-care-response)

بإمكانك مشاهدة معلومات عن كوفيد-19 في الموارد التالية:

-منظمة الصحة العالمية (WHO)

[www.who.int/emergencies/disease/novel-coronavirus-19](http://www.who.int/emergencies/disease/novel-coronavirus-19)

- مكتب منظمة الصحة العالمية الإقليمي لشرق البحر المتوسط

[www.emro.who.int/health-topics/corona-virus/information-resources.html](http://www.emro.who.int/health-topics/corona-virus/information-resources.html)

- مكتب منظمة الصحة العالمية لأوروبا

[www.euro.who.int/en/health-topics/health-topics/health-emergencies/coronavirus-covid-19](http://www.euro.who.int/en/health-topics/health-topics/health-emergencies/coronavirus-covid-19)

---

شكراً جزيلاً على وقتك ولمشاركتك في هذا الاستبيان

إذا كانت لديك أية أسئلة أو مخاوف، بإمكانك التواصل مع صباح بوفخيد على البريد الإلكتروني [sabah.boufkhed@kcl.ac.uk](mailto:sabah.boufkhed@kcl.ac.uk)



## Appendix III

**COVID-19 Situation in the Responding Services (n = 26):  
Description of Cases**

	n	%
Type of cases reported		
Confirmed	14	52
Confirmed + suspected	3	11
Confirmed + suspected + probable	2	7
Confirmed + probable	1	4
Suspected	6	22
Probable	1	4
Cases reported among		
Patient	8	30
Patient + relative + physician + nurse <sup>a</sup>	4	15
Patient + nurse	3	11
Patient + relative	2	7
Patient + relative + nurse <sup>b</sup>	2	7
Patient + physician + nurse	2	7
Patient + physician	1	4
Physician	2	7
Nurse	2	7
Missing	1	4
Location of the cases identified		
In the service	9	33
Another service of the hospital where the palliative care is located	15	56
Both in the service and another service	2	7

<sup>a</sup>One respondent also specified Other: "Administrative staff".

<sup>b</sup>Two respondents also specified "Other": "manager, coordinator"; "cleaning staff, secretary, kitchen staff".

## Appendix IV

**COVID-19 Situation in the Responding Services (n = 26):  
Case Identification and Actions Taken by the Service**

	n
Case identification	
Phone call	10
Hospital dashboard/hospital HIS/infectious diseases teams	8
COVID-19 screening of patients at admission and with symptoms; and of health-care providers	3
Symptoms identified	2
Diagnosed done by service doctor	1
Department officer	1
Call for support	1
Routine examination	1
Laboratory report	1
Missing	2
Actions taken	
Referral	14
Isolation/containment measures	10
Reporting	9
Communication with staff	6
Treatment	2
Protection of staff	1
Communication with users	1
Infection control involvement	1
Communication with department head	1
COVID team managed case	1
Home quarantine (for infected staff and contacts)	1
Training with staff	1
Contact tracing	1
Testing of all staff	1
Asymptomatic cases were followed up as outpatient	1
Use of smartphone app to follow-up COVID patients	1
All cancer hospital entrances were closed except for 1 inpatient and 1 outpatient entrance with triage with symptoms screening	1
All patients and caregivers to wear a mask (offered if do not have one)	1
Informed staff of SOP for COVID protection and case detection	1
Daily update and assessment meeting in the unit	1
Missing	3

*Appendix V*  
**Mechanisms in Place to Communicate and Coordinate  
the Response (N = 43)**

	<i>n</i>	<i>%</i>
Receiving information		
Institutions or person who would inform the hospice/service <sup>a,b</sup>		
Infection control team	13	30
Ministry of Health (MoH)/provincial health directorate	7	16
Head of hospital and/or department/hospital management/administration	6	14
Medical staff (doctors and/or nurses)	3	7
Laboratory/laboratory review online	2	5
Medical/professional society	1	2
Preventive medicine team	1	2
Emergency service	1	2
Do not know or N/A or missing	10	23
Person who would be informed in the hospice or service <sup>a,b</sup>		
Designated doctor/doctor in chief/medical director/nursing manager	8	19
Head of hospital and/or department/hospital management	7	16
Infection control team	5	12
Medical/clinical chief	4	9
Medical staff (doctors and/or nurses)	4	9
COVID-19 team	3	7
All staff	2	5
Specialist palliative care physician and consultant/consultant in charge of patient	2	5
MoH	1	2
Professor	1	2
Do not know or missing	10	23
Communication system(s) that will be used to receive information: <sup>b</sup>		
Mobile phone available 24/7	33	77
WhatsApp/Viber group	21	49
Telephone (in the service)	21	49
Email	16	37
Focal point person identified in the service responsible for collecting and sharing up-to-date information		
Yes <sup>c</sup>	31	72
No	7	16
Unsure	4	9
Missing	1	2
Sharing information		
Any communication means in place to share COVID-19 or other urgent information with staff <sup>b</sup>	41	95
Phone call	28	65
WhatsApp/Viber	27	63
Email	20	47
Text message	14	33
With patients <sup>b</sup>	39	91
Text message	18	42
WhatsApp/Viber	10	23
Phone call	35	81
Email	4	9
With relatives, visitors <sup>b</sup>	39	91
Phone call	37	86
Text message	13	30
WhatsApp/Viber	10	23
Email	2	5

<sup>a</sup>Data obtained from the analysis of open-text questions.

<sup>b</sup>Multiple choices allowed.

<sup>c</sup>19 respondents provided details: medical staff in charge (physician/nurse) (*n* = 9); infection control unit (*n* = 3); head of department/service (*n* = 2); liaison officer (*n* = 1); hospital manager/executive (*n* = 1); doctors (*n* = 1); COVID-19 team (*n* = 1); administrative assistant (*n* = 1).

*Appendix VI*  
**Information Available in the Service (N = 43)**

	Paper-Based Registry <sup>a</sup>	Electronic Record <sup>a</sup>	None	Other*
	n (%)	n (%)	n (%)	n (%)
Up-to-date contact list of				
All staff working in or for the service	20 (47)	31 (72)	2 (5)	1 (2)
All patients that attended or have attended the service	17 (40)	29 (67)	2 (5)	1 (2)
All relatives that visited or have visited the service	14 (33)	13 (30)	16 (37)	1 (2)
Patients visited in the community	13 (30)	14 (33)	16 (37)	1 (2)
System collecting information about				
Patients' symptoms	21 (49)	33 (77)	0 (0)	2 (5)
Patients' outcomes	18 (42)	31 (72)	0 (0)	2 (5)
Treatment given	19 (44)	35 (81)	0 (0)	1 (2)
Dates of patients' visits or stay	19 (44)	34 (79)	1 (2)	2 (5)
Dates of relatives' visits	15 (35)	14 (33)	16 (37)	2 (5)

<sup>a</sup>Multiple choices allowed.

*Appendix VII*

**Tables Summarizing the Qualitative Analysis of Open-text  
 Questions: Challenges Foreseen in the Upcoming Month  
 (N = 30)**

	n
Factors External to the Service	10
Social distancing	2
Fast community spread/community commitment to PPE	2
Second wave	2
Increased number of cases	2
Containment of the diseases	1
Psychosocial consequences of lockdown	1
Service organization	7
Reorganization of work	1
Extension of PC to COVID 19 patients/increase in patients with COVID 19 postintensive care	2
Less patient	1
Admission control	1
Inability of patients to attend the outpatient clinic	1
Limitations in home visits	0
Repeating tests	1
Staff workload and well-being	6
Manage the increased stress and anxiety	3
Inadequate staffing	1
Increased workload	1
Staff infected or quarantined	1
Quality of care (not optimum during a pandemic, delay in treatment)	3
Resources	3
Financial concern and burden (upcoming financial crises- donations)	1
Not enough sterilization equipment	1
Inadequate medicines	1
Other	2
Infection among staff/patient	1
Health system overload	1

*Appendix VIII*

**Tables Summarizing the Qualitative Analysis of Open-text  
Questions: Respondents' Views on Help Needed  
(N = 24)**

	<i>n</i>
Infection control	8
Quick control of COVID-19/vaccine	2
Barrier measure and screening	2
Preventive measures/social distancing and hygiene	3
Isolation	1
Resources for service	6
Training/rapid training and orientation	2
Sufficient staff number	1
Getting tests	1
Financial help and support	2
Regulations	3
Obey instructions	1
Lockdown	1
Fines for violators	1
Team support	3
Team work/meeting	1
Psychosocial support	1
More support from service/administration	1
Individual behavior	2
Awareness raising	1
Individuals being careful	1

*Appendix IX*

**Tables Summarizing the Qualitative Analysis of Open-text  
Questions: Respondents' Biggest Worries (N = 22)**

	<i>n</i>
Getting infected and transmitting COVID 19	11
Getting ill/nosocomial COVID	7
infecting family and patients	4
Impact of COVID on healthcare	6
Closing service due to COVID	1
Patient outcome	1
Resources drained for COVID and negative impact on other services including PC	1
Infection control	4
Asymptomatic transmission	1
Spike in cases that would overwhelm capacity/second wave/not being able to control the virus	3
Impact of COVID on society	2
The changes after pandemic	1
Related socioeconomic problems and medical problems	1
Other	1
To forget	1

*Appendix X***Tables Summarizing the Qualitative Analysis of Open-text Questions: Limitations to Share Expertise (N = 10)**

	<i>n</i>
Service overload	3
Staff shortage	1
Time restraints	1
Work pressure	1
Training/awareness	2
Lack of education of HCP/lack of knowledge of PC role	1
Lack of training	1
Lack of integration of PC into oncology	1
Attitudes of HCP	1
Communication problems	1
Most of the resources and efforts directed to COVID response	1
Limit of consultation because of the use of video conferencing	1
Single center experience	1

*Appendix XI***Tables Summarizing the Qualitative Analysis of Open-text Questions: Services That Could Be Provided Remotely (N = 33)**

	<i>n</i>
Nonmedical palliative care	34
Psychological	17
Social	2
Spiritual	7
Bereavement and grief support	7
Nutrition	1
Medical care/consultations	14
Consultations/Medical support and care	5
Follow-up of patients and family	1
Pain and symptom management	6
Treatment/medication refill	2
End-of-life management	9
Managing end of life	8
Communication with family at the end of life	1
Education	3
Caregiver education	2
Teaching	1
Other	2
Communication with HCW	1
COVID-19 positive consultation/referrals	1

## Appendix XII

**Tables Summarizing the Qualitative Analysis of Open-text Questions: Disadvantages and Advantages of Using Technology for Providing Palliative Care (N = 27)**

	<i>n</i>
Disadvantages of using technology	
Resources	9
Internet connection/accessibility	4
Lack of technology devices for some patients/society	4
Time pressure	1
Trust and cooperation	4
Cooperation from the patients' family/relatives hiding information from patients	2
Difficult to build rapport/trust	2
Appropriateness issues	3
Difficult for elderly patients	1
Difficult for end-of-life care	1
Difficulty or lack of knowledge to use technology	1
Lack of body language when not face-to-face communication	3
Difficulty in documenting, medical evaluation or examining patient, and assessing symptoms	2
Acceptance issues	2
Adaptation	1
Acceptance from society	1
None	2
Other	3
Psychosocial issues	1
Difficulty in reaching individuals	1
Inaccuracy in some appointments and related issues in session program	1
Advantages of using technology	
Remote care delivery and management	8
Medical care, pain management	2
Appointment and follow-up	2
Communication between medical care personnel and patients	2
To postpone follow-up	1
Relief and psychological support for patients to face the crisis	1
Mean to control transmission (less risky/enabled to protect ourselves/stay confined at home)	6
Communication - generic	2
Communication	1
Communication with HCP	1
Other	7
Support	2
Working well	2
Convenient, easy, practical	1
Direct contact	1
Saves time and effort	1
Missing	6

HCP = health-care provider.